



## Conservation Reserve Program

### ESTABLISHING AND MAINTAINING TREES AND SHRUBS

Natural Resources Conservation Service (NRCS)

October 2003

#### INTRODUCTION

Trees and shrubs may be planted to reduce soil erosion and sedimentation, improve water quality, and provide wildlife habitat.

This job sheet provides instructions for planting and maintaining trees and shrubs so that they can serve their intended purpose. Using proper planting and management techniques, especially during the establishment years, will significantly improve plant health and survival.

#### SITE PREPARATION

Before planting, it is essential to reduce competition from other vegetation that may be present on the planting site, such as grasses or weeds. The type and density of existing vegetation will determine how much pre-planting control is needed.

It's important to allow adequate time to complete this process. If significant amounts of noxious or aggressive weeds or invasive plants are present, be aware that you may need a year or two to control them before you can plant, especially if you will be planting a large area. By state law, noxious weeds in Maryland are Johnsongrass, shattercane, Canada thistle, bull thistle, plumeless thistle, and musk thistle.

For more information about controlling specific weeds in tree and shrub plantings, contact your local office of Maryland Cooperative Extension; the Maryland Department of Agriculture, Weed Control Section; or the Maryland Department of Natural Resources, Forest Service.

#### Sites without Existing Vegetation

If the seedling trees and shrubs will be planted into a clean, relatively weed-free area (such as cropland that was planted during the previous growing season), then competition from existing vegetation should not be a concern. However, a companion planting (see page 2) may be needed or desirable for erosion control and/or to reduce future weed competition.

Take into account any noxious or aggressive weeds on the site that might have been suppressed (but not killed) with previous herbicide applications. If live rootstocks are present, these weeds may be very



difficult to kill in a new planting without destroying the desirable plants. If you think you have a significant weed problem, it may be prudent to plant a temporary cover crop such as clover or spring oats and use an appropriate herbicide to treat weeds for one full growing season. Then plant the trees and shrubs the following year. If you don't know the site's weed history, consider contacting the local Maryland Department of Agriculture weed control specialist. The local specialist may have a record of weed control assistance previously provided on the property.

#### Sites with Existing Vegetation

If trees and shrubs are going to be planted into existing vegetation (for example, grasses, weeds), you will need to reduce competition before planting. For sites that need extensive preparation, much of the work can be done during the fall prior to spring planting.

Mow or brush hog the field or planting strips. Then either treat the area with an appropriate herbicide or cultivate the planting area to reduce competition.

**Using herbicides.** Choose a non-selective herbicide such as glyphosate (for example, Roundup, KleenUp). A selective herbicide such as 2,4-D may be used instead, depending on the species of weeds you are trying to control. Follow all label directions when using herbicides, and consider herbicide persistence (carryover) as it may affect new plantings.

For extremely vigorous turf or weeds, you should plan to make one application of herbicide in early fall, followed by another the next spring before planting. Or, if you make the first herbicide application in the spring, you should plan to make a second application a few weeks before planting, if needed.

Do not plant seedling trees and shrubs until the competing vegetation is sufficiently controlled. It is much easier to control the competition before planting than afterward. Cultivation of the planting area may be needed following herbicide treatment if the dead plant matter is very thick and will be difficult to plant through. You may also need to re-spray after cultivation if weed seeds brought to the surface germinate.

**Using cultivation only.** If you do not want to use herbicides, then you will need to cultivate the field or planting strips. Cultivation is usually less effective than herbicides for killing heavy sod or persistent weeds. Also, the bare ground produced by cultivation may be subject to erosion and can provide a good seedbed for more weed growth. If necessary, use a companion planting mix (as described below) to control erosion and suppress weeds.

**Companion Planting**

A companion planting mix of herbaceous species may be needed or desirable for erosion control and/or to reduce future weed competition, especially on large cultivated areas or where mulching is not feasible. Be aware that if you don't provide ground cover, then nature will. Nature's choice is likely to be weeds that you will need to control.

The following mix of three fine fescue grasses, plus clover, is recommended:

*Table 1. Recommended companion seeding mix for tree and shrub plantings.*

Fine Fescue & Clover Mixture	Seeding Rate (lbs/ac)
Creeping Red Fescue or Chewings Fescue	3 - 6
Hard Fescue	3 - 6
Sheep Fescue	3 - 6
White Clover	1 - 2

Alternatively, planting only the fine fescues (no clover), or only one fine fescue species plus clover, is also acceptable. The total seeding rate for the grasses should be at least 9 pounds per acre, up to 18 pounds per acre. Use the higher rate for erosion control, and the lower rate if only weed suppression is needed. The

companion mix can be planted in the fall prior to a spring planting of trees and shrubs, or in the spring along with the tree/shrub planting.

**Herbicide Carryover**

Carryover from herbicide treatments (recently applied or from prior years) can pose a threat to new plantings. Seedlings are particularly sensitive to herbicide carryover. Herbicides such as Basagran, Blazer, Poast, Plateau and Roundup have low persistence and generally do not pose a risk for carryover. Herbicides such as Atrazine, Preview, Canopy, Classic, Lorox Plus, Command, Scepter and Treflan have medium to high persistence and can pose a risk of carryover. The persistence of herbicides is directly affected by factors such as soil pH and moisture. To assess risks before planting, read the herbicide label or contact the manufacturer for specific information on persistence.

**PLANTING TREES AND SHRUBS**

Trees and shrubs that are planted correctly will grow faster, and will be more likely to survive, than ones that are planted incorrectly.

Always check for utility lines (water, gas, electric, cable) before planting. Avoid planting on top of buried utility lines or below overhead lines. When planting along property lines, consider using a 6 to 10-foot setback for the first row of seedlings to avoid potential conflicts with neighbors. If the planting is adjacent to cropland, consider using small trees or shrubs in one or two rows next to the field to reduce future problems with overhanging branches.

**Plant Availability and Planting Dates**

Bare-root plants are typically available only during late winter to early spring, and generally must be planted during that time. To obtain recommended planting dates for your area, contact your local NRCS Field Service Center.

**Storing and Planting Techniques**

Bare-root seedlings must be properly handled and stored to ensure a successful planting. When the seedlings arrive, open the nursery bales and check the plants for overheating, molding, or dryness. If they are satisfactory, store them in a cool, shady place with the roots protected, and plant within a few days.

If the seedlings cannot be planted immediately, dig a trench about a foot deep and bury the seedlings' roots in soil. This is called "heeling-in." Pack the soil firmly, water thoroughly, and make certain all roots are covered. The seedlings may also be stored well wrapped in a refrigerator as long as the plants are still dormant.

Evergreens require very careful handling. When heeling-in evergreens, split the small bundles and spread out the seedlings in a trench to make sure the root system of each seedling is protected by soil.

Bare-root seedlings can be machine planted, or hand planted with a planting bar or dibble. See Figure 1 for hand planting information.




Figure 1. Hand planting bare-root tree and shrub seedlings using a planting bar or dibble.

### Lime and Fertilizer

Newly planted trees and shrubs should not be limed and fertilized, unless soil tests show that pH and

nutrients are extremely low. For most sites, it's best to allow the root systems of new plantings to become established before applying lime and fertilizer.

### PROTECTING PLANTS

Plants should be protected from damage by wildlife, human activities, or livestock by using rodent guards, repellents, tree shelters, fences, and/or other exclusion measures.

#### Rabbit and Rodent Damage

Rabbits and rodents can use tall grasses and weeds for protection while gnawing on tree and shrub plantings. Mow, mulch, or use herbicides as needed to control vegetation around plantings. Follow recommendations from Maryland Cooperative Extension when using repellents or poisons to protect plantings from mice and voles.

#### Tree Shelters

Tree shelters may be used to protect seedlings from competition from weeds, damage by deer and small mammals, and damage by people while mowing or trimming around plants. Tree shelters should be a minimum of 3 feet tall to protect seedlings from deer damage. Use shelters that are at least 4 feet tall if the potential for deer damage is severe. Otherwise, shelters that are a minimum of 2 feet tall can be used to reduce damage caused by rodents, rabbits, or maintenance equipment. Push each shelter into the soil to a depth of at least 1 inch to exclude rodents.

Stake each shelter with a wooden stake (minimum 1-inch thickness), or a plastic or fiberglass post, that is at least the same height as the tree shelter being used. Do not use metal or bamboo stakes. Install and maintain bird exclusion netting on the tops of tree shelters until the plantings extend out of the tubes.

#### Fences and Use Exclusion

Many types of fences and exclusion devices are available. Contact your local NRCS Field Service Center for recommendations for your site.

### ESTABLISHING THE PLANTING

#### Planting Year

Control weeds by mowing or treating with an appropriate herbicide. Nesting season restrictions on disturbance of the planting do not apply during the establishment period. Mowing should be done with extreme caution to avoid damaging the stems or bark of plantings, especially on seedlings. Vegetation not immediately adjacent to the plantings, such as between rows, can be left for wildlife food and cover (except for

## CRP - Trees and Shrubs (Bare-Root Seedlings) Job Sheet - 4

noxious weeds, which must be controlled as required by state law). If you need assistance identifying noxious weeds, contact your local NRCS Field Service Center; Maryland Cooperative Extension; or Maryland Department of Agriculture, Weed Control Section.

Herbicides may be spot-sprayed around plantings or applied to the planting strip. Follow specific label instructions to reduce or eliminate damage to trees and shrubs. Do not apply herbicides on windy days when spray drift can damage nearby plantings.

If tree shelters are used, periodically check to make sure that shelters are upright and the bird exclusion netting is in place.

### Second Year After Planting

Continue to control weeds by mowing or treating with an herbicide. Always avoid damaging the plantings during mowing and herbicide application.

Continue to inspect tree shelters and make sure they are upright and functioning as intended.

### MAINTAINING ESTABLISHED PLANTINGS

Trees and shrubs should be well established by the third year after planting.

CRP participants must maintain enrolled practices for the life of the contract. "Maintenance" refers to activities that are carried out as needed to keep plantings in good condition so they will continue to function as planned.

Control noxious weeds and other invasive plants by spot treatment, using mechanical methods or approved herbicides. If it becomes necessary to control noxious weeds during the nesting season, contact your local weed control specialist concerning recommendations for spot-treating the weed problem. Spot treatment is limited to the immediate area of infestation.

In an established planting, you must request and receive approval from the FSA County Committee before spraying or mowing during the nesting season. For more information about controlling specific weeds, contact your local office of Maryland Cooperative Extension; the Maryland Department of Agriculture, Weed Control Section; or the Maryland Department of Natural Resources, Forest Service.

Periodic mowing for cosmetic purposes is prohibited at all times, and annual mowing for generic weed control is also prohibited. If natural regeneration of native trees and shrubs occurs, do not remove them or suppress them with herbicides or other methods. Allow a natural forested community to develop.

If tree shelters were used, remove them two years after the trees grow out of the shelters or when the trees are 2-inch caliper (as measured at 6 inches above the ground). It may be more convenient to stagger the removal of the shelters over a period of 2 to 3 years. All shelters should be removed by year 7.

*Please note: Brand names are mentioned in this jobsheet for informational purposes only. NRCS does not intend any endorsement of brands mentioned, nor criticism of similar products not mentioned.*

*Contents of this jobsheet may be reproduced for non-commercial purposes, provided that USDA-NRCS, Maryland, is credited. Text, graphics, and photo by Anne Lynn, State Biologist, NRCS, Maryland. (Portions adapted from a jobsheet prepared by Livy Marques, formerly with NRCS, Maryland.)*

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, sexual orientation, and marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call (202) 720-5964 (voice or TDD). USDA is an equal opportunity provider and employer.

